

MICROCELL HYBRID CLONES

CHROMOSOME	C8 and others from group c										N2 and others from group N				I
	B-5	B-24	D-1	D-2	D-8	D-14	E-21	I-23	L-34	M-16	T	W-23	Y-68	Z-15	
S207	X	X	X	X	●	●	●	●	X	X	X	X	X		
S177	X	X	X	X	●	●	●	●	●	X	●	X	X		
Vk1-2	●	●	X	X	●	●	●	●	●	X	X	●	X		
Vk3-2	●	●	●	●	●	●	●	●	●	●	●	●			
Ck	●	●	X	X	●	●	●	●	●	X	●	●			
S156	X	X	X	X	●	●	●	●	●	X	X	X	X		
S159	X	X	X	X	●	●	●	●	●	X	X	X	X		
	I	I	T	I	I	I	I	I	●	X	T	I			I

I: INDEPENDENT T: TRANSLOCATED

FIG. 1




FIG. 2





FIG. 3

B : BRAIN
 L : LIVER
 SM : SKELETAL MUSCLE
 H : HEART
 Sp : SPLEEN
 Th : THYMUS
 Ov : OVARY
 K : KIDNEY
 nc : NEGATIVE CONTROL
 pc : POSITIVE CONTROL



DIA1 : HUMAN CYTOCHROME b5 REDUCTASE

FIG. 4




FIG. 5

09/763362

B : BRAIN
H : HEART
Th : THYMUS
L : LIVER
Sp : SPLEEN
K : KIDNEY
Ov : OVARY
SM : SKELETAL MUSCLE
M : MOLECULAR WEIGHT MARKER



MB : HUMAN MYOGLOBIN

DIA1 : HUMAN CYTOCHROME b5 REDUCTASE

FIG. 6




FIG. 7




FIG. 8




FIG. 9





FIG. 10

B : BRAIN
K : KIDNEY
Sp : SPLEEN
H : HEART
L : LIVER
Th : THYMUS
pc : POSITIVE CONTROL
nc : NEGATIVE CONTROL
M : MOLECULAR WEIGHT MARKER



IgM : HUMAN IMMUNOGLOBULIN μ CHAIN

FIG. 11




FIG. 12

FIG. 13




FIG. 14





FIG. 15






FIG. 16

FIG. 17




FIG. 18




FIG. 19




FIG. 20





FIG. 21






FIG. 22

FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27




FIG. 28




FIG. 29 TRANSFORMANT TT2F CELL CLONES




FIG. 30





FIG. 31



RESTRICTION ENZYMES:

Kp: KpnI, B: BamHI, Bg2: Bgl2, RI: EcoRI, RV: EcoRV, N: Not, SI: SalI, Sca: ScaI, Sfi: SfiI,
 Sm: Sma, X: XhoI, (X): XhoI RESTRICTION SITE FROM λ VECTOR
 dK: KpnI RESTRICTION SITE DELETED, dX: XhoI RESTRICTION SITE DELETED
 (Sm/SI): LIGATED TO SmaI RESTRICTION SITE AFTER SalI BLUNTING
 (SI/RV): LIGATED TO EcoRV RESTRICTION SITE AFTER SalI BLUNTING
 DOTTED PORTION: pBluescript SKII(+) OR pUC18 PLASMID-DNA

◁▷ : LoxP SEQUENCE

FIG. 32
HIGH CONCENTRATION G418 RESISTANT TT2F CELL CLONES





FIG. 33



CHROMOSOMES

Inventor(s): Kazuma TOMIZUKA et
al.


DOCKET NO.: 081356/0158

09/763362



FIG. 34

FIG. 35






FIG. 36




FIG. 37




FIG. 39

	BRAIN		SPLEEN		LIVER		BONE MARROW		TESTIS				
									1ST MEIOSIS		2ND MEIOSIS		SPERM
									+	-	+	-	
16-5 (F1)	30 (100%)	0	11 (21%)	42	30 (97%)	1	30	15 (100%)	15 (100%)	0	15 (94%)	1	11 (32%)
17-8 (F1)	30 (100%)	0	12 (30%)	28	30 (100%)	21 (43%)	28	16 (100%)	16 (100%)	0	9 (53%)	7	14 (34%)
17-23 (F1)	41 (95%)	2	7 (17%)	34	31 (97%)	5 (17%)	25	15 (100%)	15 (100%)	0	13 (81%)	3	13 (52%)

RETENTION OF SC20 FRAGMENT IN MICE





FIG. 40



b) CONTROL

a) CLONE #28

FIG. 41

	HK23	HK28	HK29
ANTIBODY HEAVY-CHAIN KNOCKOUT	HOMO	WILD-TYPE OR HETERO	WILD-TYPE OR HETERO
ANTIBODY LIGHT-CHAIN KNOCKOUT	HETERO	WILD-TYPE	HETERO
W23 FRAGMENT	+	+	+
SC20 FRAGMENT	+	+	+
HUMAN μ CHAIN IN SERUM (mg/l)	100	5.9	14
HUMAN κ CHAIN IN SERUM (mg/l)	8.6	8.4	25
HUMAN μ/κ CHAINS IN SERUM (mg/l)	18	0.13	BELOW DETECTION LIMIT

* SINCE "ANTIBODY HEAVY-CHAIN KNOCKOUT" IS JUDGED BY THE PRESENCE OR ABSENCE OF THE EXPRESSION OF MOUSE μ CHAIN, IT IS IMPOSSIBLE TO DISCRIMINATE HETERO FROM WILD-TYPE.

* "ANTIBODY LIGHT-CHAIN KNOCKOUT" IS JUDGED BY SOUTHERN BLOT ANALYSIS.

FIG. 42





FIG. 44






FIG. 45




FIG. 46




FIG. 47




FIG. 48





FIG. 49




FIG. 50




FIG. 51






FIG. 52



$\Delta H/\Delta H$, SC20: Single Tc/KO mouse

$\Delta H/\Delta H$: antibody heavy-chain KO mouse

FIG. 53



09/763362




FIG. 55




FIG. 56




FIG. 57

FIG. 58




FIG. 59





FIG. 60






FIG. 61




FIG. 62




FIG. 63





FIG. 64

FIG. 65








FIG. 66

FIG. 67





Human chromosome 2

FIG. 68

FIG. 69




FIG. 70

pBS185hisD





FIG. 71



09/763362

09/763362




FIG. 72

FIG. 73




FIG. 74




FIG. 75

